

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/70	A2	(11) International Publication Number: WO 98/58086 (43) International Publication Date: 23 December 1998 (23.12.98)
(21) International Application Number: PCT/US98/11652 (22) International Filing Date: 4 June 1998 (04.06.98) (30) Priority Data: 08/876,546 16 June 1997 (16.06.97) US (71) Applicant: ABBOTT LABORATORIES [US/US]; Chad 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: KROEGER, Paul, E.; 2416 Deerpath Drive, Lindenhurst, IL 60046 (US). ABRAYAYA, Klara; 606 Leanington, Wilmette, IL 60091 (US). BSPING, Claudia, A., Cygan; 321 Crissey Avenue, Geneva, IL 60134 (US). GORZOWSKI, Jacek, J.; 210 N. Linden Drive, Round Lake Park, IL 60073 (US). HOENLE, Robert, J.; 97 Timber Hill Drive, Crystal Lake, IL 60014 (US). MOORE, Jennifer, J.; Apartment 1, 2520 N. Seminary, Chicago, IL 60614 (US). (74) Agents: YASGER, Paul, D. et al.; Abbott Laboratories, Chad 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEIC ACID PRIMERS AND PROBES FOR DETECTING HIV-1 AND HIV-2 (57) Abstract provided herein are primer/probe sets useful for detecting HIV (i.e. either or both HIV 1 or HIV 2) in a test sample. The primer/probe sets can be employed according to nucleic acid amplification procedures including PCR or RT PCR. The primer/probe sets can also be provided in the form of a kit with other reagents for performing a nucleic acid amplification reaction.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NUCLEIC ACID PRIMERS AND PROBES FOR DETECTING HIV-1 and HIV-2

Field of the Invention

The present invention relates to HIV. In particular the invention relates to oligonucleotides and methods for detecting HIV 1 and HIV 2.

5 Background of the Invention

Viruses classified as HIV contain RNA as their genetic information and the infectivity of HIV depends upon the virus's ability to insert its genetic information into the DNA of a host. In order to insert its genetic information and therefore successfully infect a host,
10 an HIV virus must convert its genetic material (RNA) into DNA so that the HIV genetic information is compatible with that of the host.

Apparently, HIV is successful at converting its RNA into DNA, given the prevalence of AIDS. However, while the virus may successfully convert RNA into DNA, the conversion is seldom accurate. In other words, the
15 DNA copy of the viral RNA is not always exact and the DNA copy can diverge from the viral RNA by several base pairs. Hence, while a host initially may be infected with a single virus particle, after several rounds of replication, the host may be infected with a genetically diverse population of viruses.

20 Although HIV is not uniformly classified, it is generally accepted that HIV 1 and HIV 2 are different viruses. Within each of these viral classifications are several groups or subtypes. For example, within the classification HIV 1, there is "group M", which is further classified into subtypes A-F, and "group O". HIV 2, on the other hand, contains subtypes
25 A-E. Subtypes of HIV 1 and HIV 2 are broken down even further into categories to numerous to mention in this context. However, it is worth mentioning that all of these divisions are based upon the genetic variance between the viruses and, according to taxonomic theory, many of these viruses are the progeny of a single virus. Hence, the numerous
30 HIV types and subtypes demonstrate the highly mutable nature of HIV and the genetic variability of the HIV genome.

The genetic variability of the virus can be attributed to the inefficiency with which the virus converts its RNA into DNA, as

mentioned above. Another theory concerning the genetic variability of the virus is that hosts can be infected with multiple different populations of HIV (which as mentioned above, can arise out of an infection by a single virus) and through the course of replication and packaging of the viral genetic information, pieces of one viral genome can be recombined with pieces of another viral genome. Hence, upon packaging of the recombined genome, a genetically distinct virus is formed. Regardless of the manner by which the virus mutates, it is clear that viruses known as HIV have genomes that are highly mutable and are therefore constantly changing. This presents those searching for methods of detecting the virus based upon its genetic information with a constantly moving target.

Although it is known that certain regions of the HIV genome are conserved, this is not to say that these regions are immune from mutation particularly if mutations in these regions do not effect the structure of a protein encoded by these regions. Hence, developing reagents and methods for detecting HIV based upon its genetic information is a continuing challenge.

Summary of the invention

The present invention provides reagents useful for detecting HIV (that is the various subtypes of HIV 1 and HIV 2) based upon the genetic information of these viruses. In particular, the reagents are in the form of primer and probe sets which can be employed according to nucleic acid amplification procedures to specifically and sensitively detect various subtypes of HIV 1 and HIV 2. Preferably, the primer/probe sets herein provided comprise two primer sequences and one probe sequence and are employed according to a reverse transcriptase (RT) PCR format.

Primer/probe sets of the invention which can be employed to detect HIV 1 are designated herein as primer/probe set 1 (SEQ. ID. NOs. 2, 3 and 4); primer/probe set 2 (SEQ. ID. NOs. 5, 6 and 7); primer/probe set 3 (SEQ. ID. NOs. 8, 9, 10 and 11); and primer/probe set 4 (SEQ. ID. NOs. 12, 13 and 14). Primer/probe sets which can be employed to detect HIV 2 are designated herein as primer/probe set 5 (SEQ. ID. NOs. 16, 17 and 18); primer/probe set 6 (SEQ. ID. NOs. 19, 20 and 21 or 22);

primer/probe set 7 (SEQ. ID. NOs. 23, 20 and 21); and primer/probe set 8 (SEQ. ID. NOs. 16, 24 and 18).

The method for detecting HIV will generally comprise the steps of
(a) forming a reaction mixture comprising nucleic acid amplification
5 reagents, at least one primer/probe set mentioned above, and a test
sample containing an HIV target sequence; (b) subjecting the mixture to
amplification conditions to generate at least one copy of a nucleic acid
sequence complementary to the target sequence; (c) hybridizing the
10 probe to the nucleic acid sequence complementary to the target
sequence, so as to form a hybrid comprising the probe and the nucleic
acid sequence complementary to the target sequence; and (d) detecting
the hybrid as an indication of the presence of HIV in the test sample.

The preferred RT PCR format will comprise the same steps as
mentioned above but the amplification reagents will comprise an
15 enzyme having reverse transcriptase activity. In addition, according to
any of the methods provided herein, step (b) can be repeated multiple
times to increase the number of target sequence copies. It will be
understood by those skilled in the art that step (b) can be repeated
through thermal cycling the reaction mixture.

Detailed Description of the Invention

The primer/probe sets provided herein comprise two primers and
at least one probe. These primer/probe sets can be employed according
to nucleic acid amplification techniques. Hence, the primers in any
25 particular primer/probe set can be employed to amplify a target
sequence. In most cases, the probe hybridizes to the copies of the
target sequence generated by one of the primers and generally
facilitates detecting any copies of the target sequence generated during
the course of the amplification reaction. All of the primer/probe sets
30 can be employed according to nucleic acid amplification procedures to
specifically and sensitively detect HIV 1 group M and group O, as well
as the various subtypes of HIV 2. Primer/probe sets for detecting HIV
1 subtypes are presented below in Table 1 and primer/probe sets for
detecting HIV 2 subtypes are presented in Table 2 below.

TABLE 1

Primer/ Probe Set	Sequence (5'-3')	SEQ. ID. NO.
1	ATTCCCTACAATCCCCAAAGTCAAGGAGT	2
	CCTGCACTGTACCCCCCAATCC	3
	ACAGCAGTACAAATGGCA	4
2	GGAGCAGAACTTTCTATGTAGATGG	5
	CATATTGTGAGTCTGTTACTATGTTTACT	6
	TAGGAAAAGCAGGATATG	7
3	GGTACAGTATTAGTAGGACCTACACCTGT	8
	GGCCATTGTTTAACTTTTGGGCCATCCA	9
	ATTAGTCCTATTGAACTGT	10
	ATAAGCCCCATAGCC	11
4	CCTAGTATAAACAATGAGACACCAGG	12
	GATCCTACATACAAGTCATCCATGTA	13
	GGATGGAAAGGATCACCA	14

TABLE 2

Primer/ Probe Set	Sequence (5'-3')	SEQ. ID. NO.
5	ACTGATGGCAGTTCATTGCATGAATTTTAAAAG	16
	TTCCACAGCTGATCTCTGCCTTCTCTG	17
	CAGAACAAGAAATACAATTC	18
6	CCTCAATTCTCTCTTTGGAAAAGACC	19
	AAATGTTGATTGGGGTATCTCCTGTC	20
	CCAAAAATAGTAGGGGG	21
	ATAGTAGCAGGAATAGA	22
7	CAATAGTAGCAGGAATAGAGTTAGG	23
	AAATGTTGATTGGGGTATCTCCTGTC	20
	CCAAAAATAGTAGGGGG	21
8	ACTGATGGCAGTTCATTGCATGAATTTTAAAAG	16
	CACAGCTGATCTCTGCCTTCTCTGTAATAGAC	24
	CAGAACAAGAAATACAATTC	18

As alluded to above, primers included in the primer/probe sets can be used to prime synthesis of copies of an HIV 1 target sequence in the case of SEQ ID NOs. 2, 3, 5, 6, 8, 9, 12, and 13; and copies of an HIV 2 target sequence in the case of SEQ ID NOs. 16, 17, 19, 20, 23 and 24. The remaining SEQ ID NOs. (SEQ ID NOs. 4, 7, 10, 11, 14, 18, 21, and 22), hybridize with the amplification products of either or both of the primer sequences found in the same primer/probe set. For example, primer/probe set 6 is specific for HIV 2 insofar as SEQ. ID. NOs. 19 and 20 prime synthesis of the HIV 2 target sequence and SEQ. ID. NOs. 21 and 22 hybridize with the amplification products produced by SEQ. ID. NOs. 19 and 20. Hence, the probe sequences are also specific for the various subtypes of HIV 1 or HIV 2.

Primer sequences generally comprise deoxyribonucleic acid (DNA), or ribonucleic acid (RNA). Probe sequences on the other hand may comprise DNA, RNA or nucleic acid analogs such as uncharged nucleic acid analogs including but not limited to peptide nucleic acids (PNAs)

which are disclosed in International Patent Application WO 92/20702 or morpholino analogs which are described in U.S. Patents Numbered 5,185,444, 5,034,506, and 5,142,047 all of which are herein incorporated by reference. Such sequences can routinely be synthesized using a variety of techniques currently available. For example, a sequence of DNA can be synthesized using conventional nucleotide phosphoramidite chemistry and the instruments available from Applied Biosystems, Inc, (Foster City, CA); DuPont, (Wilmington, DE); or Milligen, (Bedford, MA). Similarly, and when desirable, the sequences can be labeled using methodologies well known in the art such as described in U.S. Patent Applications Numbered 5,464,746; 5,424,414; and 4,948,882 all of which are herein incorporated by reference.

The term "label" as used herein means a molecule or moiety having a property or characteristic which is capable of detection. A label can be directly detectable, as with, for example, radiolabels, fluorophores, chemiluminophores, enzymes, colloidal particles, fluorescent microparticles and the like; or a label may be indirectly detectable, as with, for example, specific binding members. It will be understood that directly detectable labels may require additional components such as, for example, substrates, triggering reagents, light, and the like to enable detection of the label. When indirectly detectable labels are used, they are typically used in combination with a "conjugate". A conjugate is typically a specific binding member which has been attached or coupled to a directly detectable label. Coupling chemistries for synthesizing a conjugate are well known in the art and can include, for example, any chemical means and/or physical means that does not destroy the specific binding property of the specific binding member or the detectable property of the label. As used herein, "specific binding member" means a member of a binding pair, i.e., two different molecules where one of the molecules through, for example, chemical or physical means specifically binds to the other molecule. In addition to antigen and antibody specific binding pairs, other specific binding pairs include, but are not intended to be limited to, avidin and biotin; haptens and antibodies specific for haptens; complementary nucleotide sequences; enzyme cofactors or substrates and enzymes; and the like.

The term "test sample" as used herein, means anything suspected of containing an HIV target sequence. The test sample is or can be derived from any biological source, such as for example, blood, ocular lens fluid, cerebral spinal fluid, milk, ascites fluid, synovial fluid, peritoneal fluid, amniotic fluid, tissue, fermentation broths, cell cultures and the like. The test sample can be used (i) directly as obtained from the source or (ii) following a pre-treatment to modify the character of the sample. Thus, the test sample can be pre-treated prior to use by, for example, preparing plasma from blood, disrupting cells or viral particles, preparing liquids from solid materials, diluting viscous fluids, filtering liquids, distilling liquids, concentrating liquids, inactivating interfering components, adding reagents, purifying nucleic acids, and the like.

A "target sequence" as used herein means a nucleic acid sequence that is detected, amplified, or both amplified and detected using the primer/probe sets herein provided. Additionally, while the term target sequence is sometimes referred to as single stranded, those skilled in the art will recognize that the target sequence may actually be double stranded. Thus, in cases where the target is double stranded, primer sequences of the present invention will amplify both strands of the target sequence.

As mentioned earlier, the primer sequences of any particular primer/probe set (by themselves or with additional oligonucleotides) can be used as amplification primers according to nucleic acid amplification procedures well known in the art. Such reactions include, but are not intended to be limited to, the polymerase chain reaction (PCR) described in U.S. Patents 4,683,195 and 4,683,202, the ligase chain reaction (LCR) described in EP-A-320 308, and gap LCR (GLCR) described in U.S. Patent No. 5,427,930 all of which are herein incorporated by reference. Generically, these exemplified amplification reactions generate multiple copies of a DNA target sequence.

In accordance with the present invention, the target sequence may indeed be DNA, on the other hand in light of the RNA nature of the HIV genome, the primer/probe sets may be employed according to an "RT PCR" format which is described in U.S. Patents numbered 5,322,770 and 5,310,652 both of which are herein incorporated by reference. Briefly,

the RT PCR format provides a method of transcribing a strand of DNA from an RNA target sequence. The copied DNA strand transcribed from the RNA target is commonly referred to as "cDNA" which then can serve as a template for amplification by any of the methods mentioned above.

5 The process of generating cDNA shares many of the hybridization and extension principles surrounding other amplification methods such as PCR, but the enzyme employed should have reverse transcriptase activity. Enzymes having reverse transcriptase activity, as well as the RT PCR process, are well known and therefore don't warrant further

10 discussion. Additionally, other methods for synthesizing cDNA are also known and include commonly owned U.S. Patent Application Serial Number 08/356,287 filed February 22, 1995, which is herein incorporated by reference.

According to a preferred embodiment, the primer/probe sets are

15 employed in the "oligonucleotide hybridization PCR" (variably referred to herein as "OH PCR") amplification reaction as described in U.S. Patent Application Serial No. 08/514,704, filed August 14, 1995, that is herein incorporated by reference. Briefly, the reagents employed in the preferred method comprise at least one primer/probe set (designated

20 herein as primer/probe sets 1-8), as well as other reagents for performing an amplification reaction. "Other reagents for performing an amplification reaction" or "nucleic acid amplification reagents" include reagents which are well known and may include, but are not limited to, an enzyme having polymerase activity, enzyme cofactors

25 such as magnesium; salts; nicotinamide adenine dinucleotide (NAD); and deoxynucleotide triphosphates (dNTPs) such as for example deoxyadenine triphosphate, deoxyguanine triphosphate, deoxycytosine triphosphate and deoxythymine triphosphate.

The preferred method generally comprises the steps of (a) forming

30 a reaction mixture comprising nucleic acid amplification reagents, at least one primer/probe set of the present invention, and a test sample suspected of containing a target sequence; (b) subjecting the mixture to amplification conditions to generate at least one copy of a nucleic acid sequence complementary to the target sequence; (c) hybridizing the

35 probe to the nucleic acid sequence complementary to the target sequence, so as to form a hybrid comprising the probe and the nucleic

acid sequence complementary to the target sequence; and (d) detecting the hybrid as an indication of the presence of the target sequence (HIV) in the sample. It will be understood that step (b) of the above method can be repeated several times prior to step (c) by thermal cycling the reaction mixture between 10 and 100 times, more typically between 20 and 60 times, as is well known in the art.

Amplification conditions are defined generally as conditions which promote annealing and extension of one or more nucleic acid sequences. It is well known in the art that such annealing is dependent in a rather predictable manner on several parameters, including temperature, ionic strength, sequence length, complementarity, and G:C content of the sequences. For example, lowering the temperature in the environment of complementary nucleic acid sequences promotes annealing. For any given set of sequences, melt temperature, or T_m , can be estimated by any of several known methods. Typically, diagnostic applications utilize hybridization temperatures which are close to (i.e. within 10°C) the melt temperature. Ionic strength or "salt" concentration also impacts the melt temperature, since small cations tend to stabilize the formation of duplexes by negating the negative charge on the phosphodiester backbone. Typical salt concentrations depend on the nature and valency of the cation but are readily understood by those skilled in the art. Similarly, high G:C content and increased sequence length are also known to stabilize duplex formation because G:C pairings involve 3 hydrogen bonds where A:T pairs have just two, and because longer sequences have more hydrogen bonds holding the sequences together. Thus, a high G:C content and longer sequence lengths impact the hybridization conditions by elevating the melt temperature.

Once sequences are selected for a given diagnostic application, the G:C content and length will be known and can be accounted for in determining precisely what hybridization conditions will encompass. Since ionic strength is typically optimized for enzymatic activity, the only parameter left to vary is the temperature. Generally, the hybridization temperature is selected close to or at the T_m of the primers or probe. Thus, obtaining suitable hybridization conditions for

a particular primer/probe set is well within ordinary skill of one practicing this art.

According to the OH PCR method, it is preferable to select primers, probes and reaction conditions such that the probe sequence
5 has a lower melt temperature than the primer sequences so that upon placing the reaction mixture under amplification conditions copies of the target sequence or its complement (variably referred to as an amplicon) are produced at temperature above the T_m of the probe. After such copies are synthesized, they are denatured and the mixture is
10 cooled to enable the formation of hybrids between the probes and any copies of the target or its complement. The rate of temperature reduction from the denaturation temperature down to a temperature at which the probes will bind to single stranded copies is preferably quite rapid for example between about 8 minutes to about 15 minutes,
15 preferably less than 2 minutes. Such a rapid cooling favors hybrid formation between the copies of the target sequence and the probe rather than, for example, hybrid formation between complementary strands of the amplicon.

In cases where labels are employed to detect primer sequence
20 products, primer sequences are labeled with either a capture label or a detection label. The probe sequence is used to hybridize with the sequence generated by the primer sequence, and typically hybridizes with a sequence that does not include the primer sequence. Similarly to the primer sequence, the probe sequence is also labeled with either a
25 capture label or a detection label with the caveat that when the primer is labeled with a capture label the probe is labeled with a detection label and vice versa. Detection labels have the same definition as "labels" previously defined and "capture labels" are typically used to separate extension products, and probes associated with any such
30 products, from other amplification reactants. Specific binding members (as previously defined) are well suited for this purpose. Also, probes used according to this method are preferably blocked at their 3' ends so that they are not extended under hybridization conditions. Methods for preventing extension of a probe are well known and are a
35 matter of choice for one skilled in the art. Typically, adding a

phosphate group to the 3' end of the probe will suffice for purposes of blocking extension of the probe.

5 Upon formation of the copy sequence/probe hybrids, the differential labels (i.e. capture and detection labels) on the copy sequence and probe sequence can be used to separate and detect such hybrids. Preferably, detection is performed according to the protocols used by the commercially available Abbott LCx® instrumentation (Abbott Laboratories; Abbott Park, IL).

10 As previously discussed, the target sequence may be DNA or the target sequence may be imbedded within the HIV genome and therefore the target sequence may be in the form of RNA. In cases where the target sequence is part of the HIV genome, it is preferable to include an enzyme having reverse transcriptase activity as part of the so-called nucleic acid amplification reagents to enable production of cDNA for
15 subsequent amplification. According to this embodiment, the primer sequences also serve as primers for cDNA synthesis. Although the invention contemplates distinct steps of cDNA production and then amplification and detection of amplified cDNA sequences, it will be understood that these processes may take place simultaneously in a
20 single amplification reaction mixture.

According to another embodiment, HIV 1 and HIV 2 can be detected simultaneously in a single reaction using a combination of two primer/probe sets (i.e. one selected from the HIV 1 specific primer/probe sets and the other selected from the HIV 2 specific
25 primer/probe sets). For example, a test sample could be contacted with primer/probe sets 1 and 8 along with amplification reagents, which may or may not include an enzyme having reverse transcriptase activity, to amplify and detect the presence of HIV 1 and HIV 2 in a test sample.

30 The oligonucleotides of the present invention also can be provided as part of a kit useful for detecting HIV 1 or HIV 2. The kits comprise one or more suitable containers containing one or more primer/probe sets according to the present invention, an enzyme having polymerase activity, deoxynucleotide triphosphates and, optionally, an enzyme
35 having reverse transcriptase activity. Typically, at least one sequence bears a label, but detection is possible without this.

The following examples are provided to further illustrate the present invention and not intended to limit the invention.

Examples

5 The following examples demonstrate detection of various subtypes of HIV-1 and HIV-2 using the primer/probe sets herein provided. These DNA primers and probes comprising the primer/probe sets are identified as SEQUENCE ID NO. 2, SEQUENCE ID NO. 3, SEQUENCE ID NO. 4, SEQUENCE ID NO. 5, SEQUENCE ID NO. 6, SEQUENCE ID NO. 7, SEQUENCE ID NO. 8, SEQUENCE ID NO. 9, SEQUENCE ID NO. 10, SEQUENCE ID NO. 11, SEQUENCE ID NO. 12, SEQUENCE ID NO. 13, SEQUENCE ID NO. 14, SEQUENCE ID NO. 16, SEQUENCE ID NO. 17, SEQUENCE ID NO. 18, SEQUENCE ID NO. 19, SEQUENCE ID NO. 20, SEQUENCE ID NO. 21, SEQUENCE ID NO. 22, SEQUENCE ID NO. 23 and SEQUENCE ID NO. 24. SEQUENCE ID NO. 2, SEQUENCE ID NO. 3, SEQUENCE ID NO. 4, SEQUENCE ID NO. 5, SEQUENCE ID NO. 6, SEQUENCE ID NO. 7, SEQUENCE ID NO. 8, SEQUENCE ID NO. 9, SEQUENCE ID NO. 10, SEQUENCE ID NO. 11, SEQUENCE ID NO. 12, SEQUENCE ID NO. 13 and SEQUENCE ID NO. 14 are specific for HIV-1. A representative target sequence from HIV-1 (subtype B, strain MN) is designated herein as SEQ ID NO. 1. SEQUENCE ID NO. 16, SEQUENCE ID NO. 17, SEQUENCE ID NO. 18, SEQUENCE ID NO. 19, SEQUENCE ID NO. 20, SEQUENCE ID NO. 21, SEQUENCE ID NO. 22, SEQUENCE ID NO. 23 and SEQUENCE ID NO. 24 are specific for HIV-2. A representative target sequence from HIV-2 (subtype A, strain NIH-Z) is designated herein as SEQ ID NO. 15. All primers and probes are consensus sequences derived from 31 HIV-1 group M (subtypes A, B and D) and group O isolates and 14 HIV-2 isolates of both the A and B subtypes for the HIV-2 sequences.

25 In the following examples, SEQ ID NO. 2, SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 12 and SEQ ID NO. 13 are used as consensus amplification primers specific for the HIV-1 target sequence. SEQ ID NO. 4, SEQ ID NO. 7, SEQ ID NO. 10, SEQ ID NO. 11 and SEQ ID NO. 14 are used as consensus internal hybridization probes for the HIV-1 amplification product. SEQ ID NO. 16, SEQ ID NO. 17, SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 23 and SEQ ID NO. 24 are used as consensus amplification primers specific for the HIV-2 target sequence

and SEQ ID NO. 18, SEQ ID NO. 21 and SEQ ID NO. 22 are used as consensus internal hybridization probes for the HIV-2 amplification product.

Example 1

Preparation of HIV Primers and Probes

5 A. HIV-1 and HIV-2 Consensus Primers Consensus primers were designed to detect all known HIV-1 or HIV-2 subtypes by oligonucleotide hybridization PCR. These primers were SEQ ID NO. 2, SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 9, SEQ
10 ID NO. 12 and SEQ ID NO. 13 for HIV-1, and SEQ ID NO. 16, SEQ ID NO. 17, SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 23 and SEQ ID NO. 24 for HIV-2. Primer sequences were synthesized using standard oligonucleotide synthesis methodology and haptenated with adamantane at their 5' ends using standard cyanoethyl phosphoramidite coupling chemistry as
15 described in U.S. Patent No. 5,424,414 Incorporated herein by reference.

 B. HIV-1 and HIV-2 Consensus Probes Consensus probes were designed to hybridize with the amplified HIV target sequence by oligonucleotide hybridization. These probes were SEQ ID NO. 4, SEQ ID NO. 7, SEQ ID NO.
20 10, SEQ ID NO. 11 and SEQ ID NO. 14 for HIV-1, and SEQ ID NO. 18, SEQ ID NO. 21 and SEQ ID NO. 22 for HIV-2. Probe sequences were synthesized using standard oligonucleotide synthesis methodology and haptenated with 2 carbazoles at the 5' end using standard cyanoethyl phosphoramidite coupling chemistry as described in U.S. Patent No.
25 5,464,746 (herein incorporated by reference), and blocked with phosphate at the 3' end.

Example 2

Sensitivity of HIV-1 Detection

30 HIV-1, subtype B has been reported as the most common strain of HIV-1 found in Europe and the United States, so it was used to assess the sensitivity of the different HIV-1 consensus primer/probe sets. RNA was isolated from gradient purified HIV-1, subtype B virions (HIV-1_{MN}, Advanced Biotechnologies Inc., Columbia, MD) using the Qiagen RNA
35 extraction procedure and column methodology as described by the manufacturer (Qiagen, Frankfurt, Germany). RNA was quantitated in

viral RNA copies per milliliter and diluted into 1 ng/ μ l ribosomal RNA to levels appropriate for detection.

Four different HIV-1 consensus primer/probe sets were evaluated: set #1 used SEQ ID NOs. 2 and 3 as primers and SEQ ID NO. 4 as probe, set #2 used SEQ ID NOs. 5 and 6 as primers and SEQ ID NO. 7 as probe, set #3 used SEQ ID NOs. 8 and 9 as primers and SEQ ID NOs. 10 and 11 as probes, and set #4 used SEQ ID NOs. 12 and 13 as primers and SEQ ID NO. 14 as probe. All sequences were prepared as described in Example 1.

Dilutions of the purified HIV-1 RNA were reverse transcribed, PCR amplified and detected using the four HIV-1 consensus primer/probe sets in separate reactions. RT-PCR was performed using 50 mM Bicine, pH 8.25, 115 mM potassium acetate, 0.5 mM EDTA, 8% glycerol, 10 μ g/ml bovine serum albumin (BSA) and 0.02% sodium azide.

Recombinant *Thermus thermophilus* polymerase was used at a concentration of 5 units/reaction, with dNTPs (dATP, dGTP, dTTP and dCTP) present at a final concentration of 0.15 mM each. Primers were used at a concentration of 500 nM each, and probes at a concentration of 10 nM each. A final concentration of 2.5 mM $MnCl_2$ was used in a total reaction volume of 0.2 ml, with sample volume of 25 to 50 μ l. The negative control was composed of 100 ng of ribosomal RNA/reaction.

Reaction mixtures were reverse transcribed and amplified in a Perkin-Elmer 480 Thermal Cycler. Reaction mixtures were first incubated at 62°C for 30 minutes to reverse transcribe the RNA, followed by 30 seconds at 94°C. PCR amplification was then initiated through a touchdown or step-down protocol to aid in the stringency of the reaction in the early stages of amplification. This utilized 8 cycles as follows: 1 cycle at 70°C for 80 seconds then 94°C for 30 seconds followed by 1 cycle of 69°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 68°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 67°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 66°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 65°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 64°C for 80 seconds then 94°C for 30 seconds, followed by 1 cycle of 63°C for 80 seconds then 94°C for 30 seconds.

Further amplification was then accomplished with 35 cycles with each cycle being 62°C for 80 seconds then 94°C for 30 seconds. After the

reaction mixtures were thermal cycled, the mixtures were maintained at 97°C for 15 minutes and probe oligo hybridization was accomplished by lowering the temperature to 4°C within 2 minutes. Samples were held at 4°C until detection of reaction products.

5 Reaction products were detected on the Abbott LCx® system (available from Abbott Laboratories, Abbott Park, IL). A suspension of anti-carbazole antibody coated microparticles and an anti-adamantane antibody/alkaline phosphatase conjugate (all of which are commercially available from Abbott Laboratories, Abbott Park, IL) were used in
10 conjunction with the LCx® to capture and detect the reaction products. The enzyme substrate used was methyl-umbelliferyl phosphate (MUP), with the rate of conversion of MUP to MU (methyl-umbelliferone) measured and reported as counts/second/second (c/s/s).

15 Data from this experiment is presented in TABLE 3 and shows detection of HIV-1, subtype B at concentrations as low as 10 molecules/assay (mol/assay) using HIV-1 primer/probe sets 1, 2 and 3, and detection of 100 molecules/assay using HIV-1 primer/probe set 4.

TABLE 3

20

[HIV-1] mol/assay	Set #1 LCx® rate	Set #2 LCx® rate	Set #3 LCx® rate	Set #4 LCx® rate
10,000	2485.4	1389.6	1179.6	1459.3
1000	2323.8	1370.4	1061.1	892.6
100	1452.7	1358.2	605.2	211.9
10	490.0	835.3	440.6	73.8
0	33.0	18.7	48.1	59.7

Example 3

Sensitivity of HIV-2 Detection

25 HIV-2, subtype A, was used to assess the sensitivity of the different HIV-2 consensus primer/probe sets. RNA was isolated from gradient purified HIV-2, subtype A virions (HIV-2_{NIH2}, Advanced Biotechnologies Inc., Columbia, MD) using the Qiagen RNA extraction procedure and column methodology as described by the manufacturer (Qiagen, Frankfurt, Germany). RNA was purified from virions that were

Initially quantitated by electron microscopy. The resulting purified RNA was diluted into 1 ng/ μ l ribosomal RNA to levels appropriate for detection.

Three different HIV-2 consensus primer/probe sets were evaluated: set #5 used SEQ ID NOs. 16 and 17 as primers and SEQ ID NO. 18 as probe, set #6 used SEQ ID NOs. 19 and 20 as primers and SEQ ID NO. 22 as probe (although SEQ ID NO 21 could also have been employed as a probe alone or in combination with SEQ ID NO. 22), and set #7 used SEQ ID NOs. 20 and 23 as primers and SEQ ID NO. 21 as probe. All sequences were prepared as described in Example 1.

Reaction mixtures were prepared, reverse transcribed, amplified and detected using the three HIV-2 consensus primer/probe sets in separate reactions, as described in Example 2.

Results presented in TABLE 4 show detection of HIV-2 to concentrations as low as 1 molecule/assay using HIV-2 consensus primer/probe sets 5 and 6, and detection of 100 molecules/assay using HIV-2 primer/probe set 7.

TABLE 4

[HIV-2] mol/assay	Set #5 LCx® rate	Set #6 LCx® rate	Set #7 LCx® rate
10,000	2254.3	1427.3	501.1
1000	2310.5	1309.6	541.2
100	2290.6	1084.9	343.6
10	1967.5	750.3	75.1
1	1116.9	371.4	49.0
0	28.4	78.6	48.1

Example 4

HIV-1 Subtype Detection

The genetic divergence of HIV is the most challenging aspect to detection. There is a need for assays that accurately detect all subtypes, including the most widely divergent "O" subtype. Serum samples were obtained from Digene (Silver Spring, MD) of HIV-1

subtypes A through F, and two isolates of HIV-1, subtype O. Viral RNA was extracted and purified from each isolate using Qiagen columns and methodology as described by the manufacturer (Qiagen, Frankfurt, Germany). RNA was quantitated in viral RNA copies per milliliter and ten-fold serial dilutions were made into 1 ng/ μ l ribosomal RNA for detection. Table 5 shows the amount of RNA (molecules/assay) used per reaction at each sample dilution for each subtype.

TABLE 5

HIV-1 Subtype	RNA molecules/assay				
	Sample Dilution				
	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}
A	1737	173	17.3	1.73	0.17
B	890	89	8.9	0.89	0.09
C	247	24.7	2.47	0.25	0.03
D	20,700	2070	207	20.7	2.07
E	7170	717	71.7	7.17	0.72
F	24,000	2400	240	24	2.4
HAM O	66,460	6646	664.6	66.46	6.65
LA O	149,000	14,900	1490	149	14.9

The four different HIV-1 consensus primer/probe sets were used in separate reactions to reverse transcribe, amplify and detect the sample RNA from the various HIV-1 subtypes as described in Example 2. Subtype O was only tested with HIV-1 primer/probe set #1. Tables 6 through 9 below show the results of this testing.

Table 6

HIV-1 Subtype	HIV-1 Set #1 LCx® rate (c/s/s)				
	Sample Dilution				
	10 ⁻¹	10 ⁻²	10 ⁻³	10 ⁻⁴	10 ⁻⁵
A	1995	1530	559	45	37
B	2103	1979	750	31	27
C	740	135	133	26	24
D	1988	2042	1907	1226	528
E	2040	1954	1584	661	71
F	2027	1967	1871	1116	461
HAM O	1315	1389	1507	1224	168
LA O	1206	1413	1509	1403	650

- 5 HIV-1 primer/probe set #1 detected all HIV-1 subtypes tested in Table 6 above, including the widely divergent O subtypes, in a sensitivity range between 100 and 1 molecules/assay.

TABLE 7

HIV-1 Subtype	HIV-1 Set #2 LCx® rate (c/s/s)				
	Sample Dilution				
	10 ⁻¹	10 ⁻²	10 ⁻³	10 ⁻⁴	10 ⁻⁵
A	1218	1207	791	165	34
B	1597	1873	1544	380	20
C	569	300	*	*	*
D	1864	1866	1796	1246	636
E	931	949	856	406	61
F	1872	1838	1882	1807	1063

HIV-1 primer/probe set #2 detected all HIV-1 subtypes tested in Table 7 above with a sensitivity similar to that of HIV-1 primer/probe set #1.

TABLE 8

	HIV-1 Set #3				
	Sample Dilution				
HIV-1 Subtype	10⁻¹	10⁻²	10⁻³	10⁻⁴	10⁻⁵
A	+	+	-	-	-
B	+	-	-	-	-
C	-	-	-	-	-
D	ND	+	+/-	-	-
E	+	+	-	-	-
F	ND	+	+	+	-

(ND = not determined)

- 5 HIV-1 primer/probe set #3 detected 5 of the 6 HIV-1 subtypes tested in Table 8 above with a sensitivity of approximately 200 mol/assay.

TABLE 9

10

	HIV-1 Set #4				
	Sample Dilution				
HIV-1 Subtype	10⁻¹	10⁻²	10⁻³	10⁻⁴	10⁻⁵
A	-	-	-	-	-
B	+	-	-	-	-
C	-	-	-	-	ND
D	+	+	+	-	-
E	+	+	+/-	-	-
F	+	-	-	-	-

(ND = not determined)

15

- HIV-1 primer/probe set #4 detected 4 of the 6 HIV-1 subtypes tested in Table 9 above with a sensitivity of between 2000 to 200 mol/assay.

Example 5**Specificity of HIV-1 and HIV-2 Primer/Probe Sets**

The specificity of HIV-1 primer/probe sets #1 and #2 was assessed against RNA which was extracted and purified from HIV-1 subtype B (strains MN and 3B) and HIV-2 serum samples as in Example 4. The two different HIV-1 consensus primer/probe sets were used in separate reactions to reverse transcribe, amplify and detect 1:10 dilutions of the sample RNA as described in Example 2. As shown in Table 10, both HIV-1 primer/probe sets detect both HIV-1 RNAs but do not detect RNA from the HIV-2 serum, indicating that the HIV-1 primer/probe sets are specific for HIV-1 detection and do not cross-react and detect HIV-2.

TABLE 10

Sample RNA	HIV-1 Set #1 LCx® rate	HIV-1 Set #2 LCx® rate
HIV-1_{MN}	2317	1321
HIV-1_{3B}	2366	1395
HIV-2	109	19

Next, the specificity of HIV-2 primer/probe set #5 was assessed against RNA isolated from gradient purified HIV-1 as in Example 2. Dilutions of the purified HIV-1 RNA were reverse transcribed, PCR amplified and detected as in Example 2 using HIV-2 consensus primer/probe set #5, with HIV-1 consensus primer/probe set #1 used in separate reactions as a control. The results, shown in Table 11, indicate that HIV-2 primer/probe set #5 does not cross-react and detect HIV-1 RNA.

TABLE 11

[HIV-1] mol/assay	HIV-1 Set #1 LCx® rate	HIV-2 Set #5 LCx® rate
1,000,000	1792	16
100,000	1819	15
10,000	1812	16
1000	1823	15
100	1714	15
10	517	15
0	27	16

Example 65 Detection of HIV-1 or HIV-2 with HIV-1 and HIV-2 Primer/Probes

In order to allow both HIV-1 and HIV-2 to be tested in one reaction instead of in two separate reactions, primer/probe sets for HIV-1 and HIV-2 would have to be used in the same reaction. The feasibility of this was tested by mixing primer/probe sets for HIV-1 and HIV-2 and using them to detect either HIV-1 or HIV-2 purified RNA.

HIV-1 and HIV-2 RNA were purified and quantitated as in Examples 2 and 3 respectively. The HIV-1 primer/probe set #1 and HIV-2 primer/probe set #8, which consisted of SEQ ID NOs. 16 and 24 as primers and SEQ ID NO. 18 as probe, prepared as described in Example 1, were used either together or separately to reverse transcribe, amplify and detect the dilutions of purified HIV-1 and HIV-2 RNA as described in Example 2. Results are shown in Table 12.

TABLE 12

[HIV-1] mol/assay	HIV-1 LCx® rate	HIV-1/2 LCx® rate	[HIV-2] mol/assay	HIV-2 LCx® rate	HIV-1/2 LCx® rate
10,000	2114.5	1858.3	10,000	2094.1	2261.6
1000	1708.0	1550.2	1000	2209.2	1966.8
100	680.8	539.6	100	2216.2	1066.6
10	112.6	151.0	10	1879.8	254.9
1	ND	ND	1	805	66.8
0	50.4	74.5	0	31	90.4

(ND = not determined)

5 While there was a 1.0 to 1.5 log decrease in detection of HIV-2
when the HIV-1 and HIV-2 primer/probe sets were used in the same
reaction mixture, the sensitivity remained acceptable at approximately
10 molecules per reaction. Importantly, detection of HIV-1 was not
10 altered by the inclusion of the HIV-2 primer/probe set with the HIV-1
primer/probe set in the HIV-1 reaction mixtures, with sensitivity
remaining at approximately 10 molecules per reaction.

15 While the invention has been described in detail and with
reference to specific embodiments, it will be apparent to one skilled in
the art that various changes and modifications may be made to such
embodiments without departing from the spirit and scope of the
invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: P. Kroeger
 K. Abravaya
 J. Gorzowski
 R. Hoenle
10 C. Esping
 J. Moore
- 15 (ii) TITLE OF INVENTION: NUCLEIC ACID SEQUENCES FOR DETECTING
 HIV-1 and HIV-2
- (iii) NUMBER OF SEQUENCES: 23
- 20 (iv) CORRESPONDENCE ADDRESS:
 (A) ADDRESSEE: Abbott Laboratories
 (B) STREET: 100 Abbott Park Road
 (C) CITY: Abbott Park
 (D) STATE: Illinois
25 (E) COUNTRY: USA
 (F) ZIP: 60064-3500
- (v) COMPUTER READABLE FORM:
 (A) MEDIUM TYPE: Floppy disk
30 (B) COMPUTER: MacIntosh
 (C) OPERATING SYSTEM: System 7.0.1
 (D) SOFTWARE: Microsoft Word 5.1a
- (vi) CURRENT APPLICATION DATA:
35 (A) APPLICATION NUMBER:
 (B) FILING DATE:
 (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
40 (A) NAME: Paul D. Yasger
 (B) REGISTRATION NUMBER: 37,477
 (C) DOCKET NUMBER: 6127.US.01
- (ix) TELECOMMUNICATION INFORMATION:
45 (A) TELEPHONE: 847/938-3508
 (B) TELEFAX: 847/938-2623
 (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2348 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- 55 (ii) MOLECULE TYPE: genomic RNA (HIV-1)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

24

	ACAGTATTAG	TAGGACCTAC	ACCTGTCAAC	ATAATTGGAA	GAAATCTGTT	50
	GACTCAGCTT	GGGTGCACCT	TAAATTTTCC	CATTAGTCCT	ATTGAAACTG	100
	TACCAGTAAA	ATTAAAGCCA	GGAAATGGATG	GCCCAAAAGT	TAAACAATGG	150
	CCATTGACAG	AAGAAAAAAT	AAAAGCATT	ATAGAAATTT	GTACAGAAAT	200
5	GGAAAAGGAA	GGGAAAATTT	CAAAAATTGG	GCCTGAAAAT	CCATACAATA	250
	CTCCAGTATT	TCCCATAAAG	AAAAAAGACA	GTAATAAATG	GAGAAAAATTA	300
	GTAGATTTCA	GAGAACTTAA	TAAGAAAAC	CAAGACTTCT	GGGAAAGTTCA	350
	ATTAGGAATA	CCACATCCTG	CAGGGTTAAA	AAAGAAAAAA	TCAGTAACAG	400
	TACTGGATGT	GGGTGATGCA	TATTTTTCAG	TTCCCTTAGA	TAAAGACTTC	450
10	AGGAAGTATA	CTGCATTTAC	CATACCTAGT	ATAACAATG	AAACACCAGG	500
	GATTAGATAT	CAGTACAATG	TGCTTCCACA	GGGATGGAAA	GGATCACCAG	550
	CAATATTCCA	AAGTAGCATG	ACAAAAATCT	TAGAGCCTTT	TAGAAAACAA	600
	AATCCAGACA	TAGTTATCTA	TCAATACATG	GATGATTTGT	ATGTAGGATC	650
	TGACTTAGAA	ATAGGGCAGC	ATAGAGCAAA	AATAGAGGAA	CTGAGACGAC	700
15	ATCTGTTGAG	GTGGGGATTT	ACCACACCAG	ACAAAAACA	TCAGAAAGAA	750
	CCTCCATTCC	TTTGGATGGG	TTATGAACTC	CATCCTGATA	AATGGACAGT	800
	ACAGCCTATA	GTGCTACCAG	AAAAAGACAG	CTGGACTGTC	AATGACATAC	850
	AGAAGTTAGT	GGGAAAATTG	AATTGGGCAA	GTCAGATTTA	CGCAGGGATT	900
	AAAGTAAAGC	AATTATGTAA	ACTCCTTAGA	GGAACCAAAG	CACTAACAGA	950
20	AGTAATACCA	CTAACAGAAG	AAGCAGAGCT	AGAAGTGGCA	GAAAAACAGG	1000
	AAATTCTAAA	AGAACCAGTA	CATGGAGTGT	ATTATGACCC	ATCAAAAGAC	1050
	TTAATAGCAG	AAGTACAGAA	GCAGGGGCAA	GGCCAATGGA	CATATCAAAT	1100
	TTATCAAGAG	CCATTTAAAA	ATCTGAAAAC	AGGCAAATAT	GCAAGATGA	1150
	GGGGTGCCCA	CACATAATGAT	GTAAAACAAT	TAACAGAGGC	AGTGCAAAAA	1200
25	ATAGCCACAG	AAAGCATAGT	AATATGGGGA	AAGACTCCTA	AATTTAGACT	1250
	ACCCATACAA	AAAGAAACAT	GGGAAACATG	GTGGACAGAG	TATACGTAAG	1300
	CCACCTGGAT	TCCTGAGTGG	GAGGTGTCTA	ATACCCCTCC	CTTAGTGAAA	1350
	TTATGGTACC	AGTTAGAGAA	AGAACCCATA	GTAGGTGCAG	AAACTTTCTA	1400
	TGTAGATGGG	GCAGCTAACA	GGGAGACTAA	AAAAGGAAAA	GCAGGATATG	1450
30	TTACTAAGAG	AGGAAGACAA	AAGGTTGTCT	CCCTAACTGA	CACAACAAAT	1500
	CAGAAGACTG	AGTTACAAGC	AATTCATCTA	GCTTTGCAAG	ATTTCAGGGT	1550
	AGAAGTAAAC	ATAGTAACAG	ACTCACAATA	TGCATTAGGA	ATCATTCAAG	1600
	CACAACCAGA	TAAAAGTGAA	TCAGAGTTAG	TCAGTCAAAT	AATAGAGCAG	1650
	TTAATAAAAA	AGGAAAAGGT	CTATCTGGCA	TGGGTACCAG	CACACAAAGG	1700
35	AATTGGAGGA	AATGAACAAG	TAGATAAATP	AGTCAGTGCT	GGAATCAGGA	1750
	AAGTACTATT	TTTAGATGGA	ATAGATAAGG	CCCAAGAAGA	CCATGAGAAA	1800
	TATCACAGTA	ATTGGAGAGC	AATGGCTAGT	GACTTTAACC	TACCACCTAT	1850
	AGTAGCAAAA	GAAATAGTAG	CCAGCTGTGA	TAAATGTGAG	CTAAAAGGAG	1900
	AAGCCATGCA	TGGACAAGTA	GACTGTAGTC	CAGGAATATG	GCAACTAGAT	1950
40	TGTACACATT	TAGAAGGAAA	AGTTATCCTG	GTCAGAGTTC	ATGTAGCCAG	2000
	TGGATACATA	GAAGCAGAAG	TTATTCCAGC	AGAGACACGG	CAGGAGACAG	2050
	CATACTTTCT	CTTAAAATTA	GCAGGAAGAT	GGCCAGTAAA	AACAATACAT	2100
	ACAGACAATG	GCCCCAATTT	CACCAGTACT	ACGGTTAAGG	CCGCCTGTTG	2150
	GTGGACGGGA	ATCAAGCAGG	AATTTGGCAT	TCCCTACAAT	CCCCAAAGTC	2200
45	AAGGAGTAAT	AGAATCTATG	AATAAAGAAT	TAAAGAAAAT	TATAGGACAG	2250
	GTAAGAGATC	AGGCTGAACA	TCTTAAGAGA	GCAGTACAAA	TGGCAGTATT	2300
	CATCCACAAT	TTTAAAAGAA	AAGGGGGGAT	TGGGGGGTAC	AGTGCAGG	2348

- 50 (2) INFORMATION FOR SEQ ID NO:2:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 55 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATTCCCTACA ATCCCCAAAG TCAAGGAGT

29

25

(2) INFORMATION FOR SEQ ID NO:3:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA

 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
CCTGCACTGT ACCCCCCAAT CC 22

(2) INFORMATION FOR SEQ ID NO: 4:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA

 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
ACAGCAGTAC AAATGGCA 18

(2) INFORMATION FOR SEQ ID NO:5:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA

 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
GGAGCAGAAA CTTTCTATGT AGATGG 26

(2) INFORMATION FOR SEQ ID NO:6:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA

 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
CATATTGTGA GTCTGTTACT ATGTTTACT 29

(2) INFORMATION FOR SEQ ID NO:7:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TAGGAAAAGC AGGATATG

18

5 (2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

15 GGTACAGTAT TAGTAGGACC TACACCTGT

29

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGCCATTGTT TAACITTTGG GCCATCCA

28

(2) INFORMATION FOR SEQ ID NO:10:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATTAGTCCTA TTGAAACTGT

20

40 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

45 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

50 ATAAGCCCCA TAGCC

15

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCTAGTATAA ACAATGAGAC ACCAGG

26

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GATCCTACAT ACAAGTCATC CATGTA

26

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGATGGAAAG GATCACCA

18

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2689 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic RNA (HIV-2)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TTGCTGCACC TCAATTCTCT CTTTGGAAAA GACCAGTAGT CACAGCACAC
ATTGAGGGTC AGCCAGTAGA AGTTTGTGTA GACACAAGGG CTAACGACTC
AATACTAGCA GGAATAGAGT TAGGGAGCAA TTATAGTCCA AAAATAGTAG
GAGGAATAGG GGGATTCATA AATACCAAGG AATATAAAAA TGTAAGAAATA
GAAGTCCTAG GTAAAAGGGT AAAAGCCACC ATAATGACAG GTGATACCCC
GATCAACATT TTTGGCAGAA ATGTTCTGAC AGCTTTAGGC ATGTCATTAA
ACCTGCCAGT TGCCAAGATA GAACCAATAA AAATAATGCT AAAGCCAGGG
AAAGATGGAC CAAGACTAAA ACAATGGCCT TTAACAAAAG AAAAAATAGA
AGCACTAAAA GAAATCTGTG AAAAAATGGA AAAAGAAGGC CAGCTAGAAG
AAGCACCTCC AACTAATCCT TATAATACCC CCACATTTGC AATCAAGAAA
AAGGACAAAA ACAAATGGAG AATGCTAATA GACTTTAGGG AGCTAAACAA
GGTAACCTAG GATTTACAG AAATTCAGTT AGGAATTCCA CACCCAGCAG
GATTGGCCAA AAAGAGGAGA ATTACTGTAC TAGATGTAGG GGATGCTTAC
TTTTCATAC CACTACATGA GGACTTTAGA CAGTATACTG CATTCACTCT
ACCATCAGTA AACATGCAG AACCAAGAAA AAGATATATA TACAAAGTCC
TACCACAGGG GTGGAAAGGG TCACCAGCAA TTTTCAATA CACAATGAGG
CAGATCTTAG AACCATTCAG AAAAGCAAAC GAGGATGTCA TTATCATTTCA
GTACATGGAT GATATCTTAA TAGCCAGCGA CAGGACAGAC TTAGAATCATG
ACAAAGTGGT CCTGCAGCTA AAAGAACTTC TAAATGGACT AGGATTTTCC
ACCCACAGATG AGAAATTCCA GAAAGACCCT CCATATCGCT GGATGGGCTA

50

100

150

200

250

300

350

400

450

500

550

600

650

700

750

800

850

900

950

1000

28

	TGAATTATGG	CCAACTAAAT	GGAAGTTGCA	AAAAATACAG	CTGCCCCAAA	1050
	AAGAAGTATG	GACAGTCAAT	GACATCCAAA	AGCTAGTGGG	TGTCCTAAAT	1100
	TGGGCAGCAC	AAATCTACCC	AGGGATAAAG	ACGAAACACC	TATGTAGGCT	1150
	AATTAGAGGA	AAAATGACAC	TCACAGAAGA	GGTACAGTGG	ACAGAATTAG	1200
5	CAGAAGCAGA	GCTAGAGGAA	AACAGAAATTA	TCTTAAGCCA	GAAACAAGAA	1250
	GGACACTATT	ACCAGGAAGA	AAAAAAGTTA	GAAGCAACAG	TCCAGAAAGA	1300
	TCAAGACAAT	CAGTGGACAT	ATAAGGTACA	CCAGGGGGAG	AAAATTCTCA	1350
	AAGGTGGGAA	AATATGCAAA	GATAAAAAAT	ACCCATACCA	ACGGGTCAGA	1400
	TTGTTAGCAC	AGGTAGTTCA	AAAAATAGGA	AAAGAAGCAC	TAGTCATTTG	1450
10	GGGACGGATA	CCAAAATTTT	ACCTACCAGT	AGAGAGAGAT	ACCTGGGAGC	1500
	AGTGGTGGGA	TAACACTGG	CAAGTAACAT	GGATCCCAGA	CTGGGACTTT	1550
	GTATCTACCC	CACCACCTGG	CAGGCTAGCA	TTTAACCTGG	TAGGAGAGCC	1600
	TGTACCAGGC	GCAGAACTT	TCTACACAGA	TGGATCCTGC	AATAGGCAGT	1650
	CAAAAGAAGG	GAAAGCAGGA	TATATAACAG	ATAGAGGGAG	AGACAGGGTA	1700
15	AAAGTATTAG	AGCAAACTAC	CAATCAGCAA	GCAGAATTAG	AAGCCTTTGC	1750
	AATGGCACTA	ACAGACTCAG	GTCCAAAGC	TAATATTATA	GTAGACTCAC	1800
	AATATGTAAT	GGGGATAGTA	GCAGGCCAAC	CAACAGAGTC	AGAGAATAGA	1850
	ATAGTAAATC	AGATCATAGA	AGAAATGATA	AAGAAAGAAG	CAATCTATGT	1900
	TGCGTGGGTC	CCAGCCCACA	AAGGCATAGG	AGGAAACCAG	GAAGTAGATC	1950
20	ATTTAGTAAG	TCAGGGCAAT	AGACAAGTAT	TATTCCTAGA	GAAAATAGAA	2000
	CCCGCGCAGG	AAGAACATGA	AAAATATCAT	AGCAATATAA	AAGAACTGTC	2050
	CCATAAAATTT	GGAATACCCA	AGCTAGTGGC	AAGACAAATA	GTAAACACAT	2100
	GTGCCCATGT	GCAACAGAAA	GGGGAGGCTA	TACATGGGCA	AGTAAATGCA	2150
	GAAGTAGGCA	CTTGGCAAAT	GGACTGCACA	CATTTAGAAG	GAAAAGTCAT	2200
25	TATAGTAGCA	GTACATGTTG	CAAGTGCATT	TATAGAGGCA	GAAGTTATCC	2250
	CACAGGAATC	AGGAAGGCAA	ACAGCACTGT	TCTTACTGAA	ACTGGCCAGT	2300
	AGGTGGCCAA	TAACACACCT	GCACACAGAT	AATGGTGCCA	ACTTCACTTC	2350
	ACAGGAAGTG	AAAATGGTAG	CATGGTGGGT	AGGTATAGAA	CAAACCTTTG	2400
	GAGTGCCTTA	CAATCCACAA	AGCCAAGGAG	TAGTAGAAGC	AATGAATCAT	2450
30	CATCTAAAAA	ATCAGATAGA	CAGAATTAGA	GAGCAGGCAA	ATACAGTAGA	2500
	AACAATAGTA	TTAATGGCAG	TTCATTGCAT	GAATTTTAAA	AGAAGGGGAG	2550
	GAATAGGGGA	TATGACCCCA	GCAGAAAGAA	TAATCAATAT	GATCACCACA	2600
	GAACAAGAAA	TACAATTCCT	CCAAGCAAAA	AATTCAAAAT	TAAAAAATTT	2650
35	TCGGGTCTAT	TTCAGAGAAG	GCAGAGATCA	GCTGTGGAA		2689

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ACTGATGGCA GTTCATTGCA TGAATTTTAA AAG

33

(2) INFORMATION FOR SEQ ID NO:17:

50 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TTCCACAGCT GATCTCTGCC TTCTCTG

27

- (2) INFORMATION FOR SEQ ID NO:18:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
CAGAACAAGA AATACAATTC 20
- (2) INFORMATION FOR SEQ ID NO:19:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
CCTCAATTCT CTCTTTGGAA AAGACC 26
- (2) INFORMATION FOR SEQ ID NO:20:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
AAATGTTGAT TGGGGTATCT CCTGTC 26
- (2) INFORMATION FOR SEQ ID NO:21:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
CCAAAAATAG TAGGGGG 17
- (2) INFORMATION FOR SEQ ID NO:22:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

30

ATAGTAGCAG GAATAGA

17

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

5

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CAATAGTAGC AGGAATAGAG TTAGG

25

15

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

25

CACAGCTGAT CTCTGCCTTC TCTGTAATAG AC

32

Claims

What is claimed is:

1. A primer/probe set for detecting HIV 1 or HIV 2 selected from the primer/probe sets consisting of: primer/probe set 1 (SEQ. ID. NOs. 2, 3 and 4); primer/probe set 2 (SEQ. ID. NOs. 5, 6 and 7); primer/probe set 3 (SEQ. ID. NOs. 8, 9, 10 and 11); primer/probe set 4
5 (SEQ. ID. NOs. 12, 13 and 14); primer/probe set 5 (SEQ. ID. NOs. 16, 17 and 18); primer/probe set 6 (SEQ. ID. NOs. 19, 20 and 21 or 22); primer/probe set 7 (SEQ. ID. NOs. 23, 20 and 21); and primer/probe set 8 (SEQ. ID. NOs. 16, 24 and 18).

2. A method of detecting HIV in a test sample, said method comprises the steps of:

(a) forming a reaction mixture comprising nucleic acid amplification reagents, at least one primer/probe set of claim 1, and a
5 test sample containing an HIV target sequence;

(b) subjecting the mixture to amplification conditions to generate at least one copy of a nucleic acid sequence complementary to the target sequence;

(c) hybridizing the probe to the nucleic acid sequence
10 complementary to the target sequence, so as to form a hybrid comprising the probe and the nucleic acid sequence complementary to the target sequence; and

(d) detecting the hybrid as an indication of the presence of HIV in the test sample.

3. The method of claim 2 wherein said amplification reagents include an enzyme having reverse transcriptase activity and said method further comprises the step, after step (a) and prior to step (b), of forming a cDNA from an HIV genome.

4. The method of claim 2 wherein said at least one primer/probe set is selected from the primer/probe sets consisting of: primer/probe set 1 (SEQ. ID. NOs. 2, 3 and 4); primer/probe set 2 (SEQ. ID. NOs. 5, 6 and 7); primer/probe set 3 (SEQ. ID. NOs. 8, 9, 10 and 11); and primer/probe set 4 (SEQ. ID. NOs. 12, 13 and 14).

5. The method of claim 2 wherein said at least one primer/probe set is selected from the primer/probes sets consisting of: primer/probe set 5 (SEQ. ID. NOs. 16, 17 and 18); primer/probe set 6 (SEQ. ID. NOs. 19, 20 and 21 or 22); primer/probe set 7 (SEQ. ID. NOs. 23, 20 and 21); and primer/probe set 8 (SEQ. ID. NOs. 16, 24 and 18).

6. The method of claim 2 wherein step (b) is repeated between 10 and 100 times

7. A kit comprising:

- a) at least one primer/probe set selected from the group consisting of: primer/probe set 1 (SEQ. ID. NOs. 2, 3 and 4), primer/probe set 2 (SEQ. ID. NOs. 5, 6 and 7), primer/probe set 3 (SEQ. ID. NOs. 8, 9, 10 and 11), primer/probe set 4 (SEQ. ID. NOs. 12, 13 and 14), primer/probe set 5 (SEQ. ID. NOs. 16, 17 and 18), primer/probe set 6 (SEQ. ID. NOs. 19, 20 and 21 or 22), primer/probe set 7 (SEQ. ID. NOs. 23, 20 and 21), and primer/probe set 8 (SEQ. ID. NOs. 16, 24 and 18); and
- b) amplification reagents.

8. The kit of claim 7 wherein said amplification reagents comprise an enzyme having reverse transcriptase activity.

9. The kit of claim 7 wherein said kit comprises at least two primer/probe sets wherein

- (a) one of the at least two primer/probe sets is selected from the group consisting of primer/probe set 1 (SEQ. ID. NOs. 2, 3 and 4), primer/probe set 2 (SEQ. ID. NOs. 5, 6 and 7), primer/probe set 3 (SEQ. ID. NOs. 8, 9, 10 and 11), primer/probe set 4 (SEQ. ID. NOs. 12, 13 and 14); and

- 10 (b) one of the at least two primer/probe sets is selected from the group consisting of primer/probe set 5 (SEQ. ID. NOs. 16, 17 and 18), primer/probe set 6 (SEQ. ID. NOs. 19, 20, 21 and 22), primer/probe set 7 (SEQ. ID. NOs. 23, 20, 21 and 22), and primer/probe set 8 (SEQ. ID. NOs. 16, 24 and 18).

THIS PAGE BLANK (USPTO)